



## Product Datasheet

<b>Product Name</b>	Interferon stimulated gene 15 Human Recombinant
<b>Cata No</b>	CB501468
<b>Source</b>	<i>Escherichia Coli.</i>
<b>Synonyms</b>	ISG15, G1P2, IFI15, UCRP, Interferon-induced 17 kDa protein precursor, ISG15 Ubiquitin-like modifier.

### Description

ISG15 ubiquitin-like Modifier is conjugated to the intracellular target proteins after IFN-alpha or IFN-beta stimulation. ISG15 enzymatic pathway is partially distinct from that of ubiquitin, differing in substrate specificity and interaction with ligating enzymes. ISG15 conjugation pathway uses a dedicated E1 enzyme, but seems to converge with the ub conjugation pathway at the level of a specific E2 enzyme. ISG15 protein targets include STAT1, SERPINA 3G/SPI2A, JAK1, MAPK3/ERK1, PLCG1, EIF2AK2/PKR, MX1/MXA, and RIG-1.

Deconjugated by USP18/UBP43 shows specific chemotactic activity towards neutrophils and activates them to induce release of eosinophil chemotactic factors. ISG15 serves as a trans-acting binding factor directing the association of ligated target proteins to intermediate filaments. ISG15 plays a role in autocrine, paracrine and endocrine mechanisms, as in cell-to-cell signaling, possibly partly by inducing ifn-gamma secretion by monocytes and macrophages. ISG15 displays antiviral activity during viral infections. ISG15 in response to IFN-TAU secreted by the conceptus, ligates to and regulate proteins involved in the release of prostaglandin F2-alpha and thus prevent lysis of the corpus luteum and maintain the pregnancy.

The interferon-stimulated gene having a molecular mass of 15 kDa is an interferon regulated gene that is induced as a primary response to diverse

microbial and cell stress stimuli, and encodes the founding member of the ubiquitin-like protein family. ISG15 shares several common properties with other ubiquitin-like molecules, but its activity is strongly regulated by precise signaling pathways that have a role in innate immunity.

ISG15 Human Recombinant fused to N-terminal Calmodulin Tag (151 a.a.) produced in E.Coli is a single, non-glycosylated polypeptide chain containing a total of 308 amino acids (157 a.a. without tag) and having a molecular mass of 34 kDa.

### Physical Appearance

Sterile Filtered colorless solution.

### Purity

Greater than 98.0% as determined by:

- (a) Analysis by RP-HPLC.
- (b) Analysis by SDS-PAGE.

### Formulation

The ISG15 solution contains 20mM Tris-HCl pH-8, 100mM NaCl and 1mM DTT.

### Stability

ISG15 although stable 4°C for 4 weeks, should be stored desiccated below -18°C.

For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).

**Please prevent freeze-thaw cycles.**

**\* For Non-Clinical Research Use Only \***



# California Bioscience

83103 Avenue 48, Ste.1B #204  
Coachella, CA 92236 USA  
Phone : +1.6268339877  
Email : [info@cali-bio.com](mailto:info@cali-bio.com)

## Sequence

MADQLTEEQI AEFKEAFSLF DKDGDGTITT  
KELGTVMRSL GQNPTEAELQ DMINEVDADG  
NGTIDFPEFL TMMARKMKDT DSEEEIREAF  
RVFDKDGNGY ISAAELRHVM TNLGEKLTDE  
EVDEMIREAD IDGDGQVNYE EFVQMMTAKG  
S

## Product Datasheet

**\* For Non-Clinical Research Use Only \***